

Computer Graphics 2016

9. Splines and Curves

Hongxin Zhang

State Key Lab of CAD&CG, Zhejiang University

2016-11-28

About homework 3

- an alternative solution with WebGL
- links:
 - WebGL lessons
http://learningwebgl.com/blog/?page_id=1217
 - My simple test
<https://github.com/hongxin/PonyGL>
- Please use google's browser: chrome

classification of curves

$$y = x^2 + 5x + 3 \quad \longrightarrow \quad y = f(x)$$

(**explicit curve**)

$$(x - x_c)^2 + (y - y_c)^2 - r^2 = 0 \quad \longrightarrow \quad g(x, y) = 0$$

(**implicit curve**)

$$\begin{aligned} x &= x_c + r \cdot \cos \theta \\ y &= y_c + r \cdot \sin \theta \end{aligned} \quad \longrightarrow \quad \begin{cases} x = x(t) \\ y = y(t) \end{cases}$$


(**parametric curve**)

classification of curves

implicit curve

- planar: $f(x,y)=0$:
 $x^2+y^2-36=0$

- 3D curves



$$\begin{cases} f(x, y, z) = 0, \\ g(x, y, z) = 0. \end{cases}$$



implicit curves

advantage of implicit curve:

To a point (x,y) , it is easy to detect whether $f(x,y)$ is >0 , <0 or $=0$.


disadvantage of implicit curve:

To a curve $f(x,y)=0$, it is difficult to find the point on it..






implicit curves

Display of implicit curves---chain coding



implicit curves

Display of implicit curves---subdivision




Parametric curves

- variable is a scalar, and function is a vector:

$$C=C(u)=[x(u), y(u), z(u)],$$

- Every element of the vector is a function of the variable(the parameter)




Parametric curves

given a curve $\mathbf{C}(u)$, its tangent is $\mathbf{T}=\mathbf{C}'(u)$.

difference of arc length:


$$(ds)^2=(dx)^2+(dy)^2+(dz)^2=((x')^2+(y')^2+(z')^2)d^2u$$

- Arc length: $s = \int_{u_0}^u ds = \int_{u_0}^u \sqrt{(x')^2 + (y')^2 + (z')^2} du$



Parametric curves and splines

- Cubic Hermite interpolation
- Catmull-Rom interpolation
- Bezier curves




Cubic Hermite interpolation

Goal: Interpolate Values




Nearest Neighbor Interpolation



Problem: values not continuous

Linear Interpolation




Problem: derivatives not continuous

Smooth Interpolation?



Cubic Hermite Interpolation



Given: values and derivatives at 2 points

Cubic Polynomial Interpolation

Assume cubic polynomial

$$P(t) = at^3 + bt^2 + ct + d$$

Why? 4 constraints => need 4 degrees of freedom

Cubic Hermite Interpolation

Assume cubic polynomial

$$P(t) = at^3 + bt^2 + ct + d$$

$$P'(t) = 3at^2 + 2bt + c$$

Solve for coefficients:

$$P(0) = h_0 = d$$

$$P(1) = h_1 = a + b + c + d$$

$$P'(0) = h_2 = c$$

$$P'(1) = h_3 = 3a + 2b + c$$

Matrix Representation

$$h_0 = d$$

$$h_1 = a + b + c + d$$

$$h_2 = c$$

$$h_3 = 3a + 2b + c$$

$$\begin{bmatrix} h_0 \\ h_1 \\ h_2 \\ h_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

Matrix Representation of Polynomials

$$P(t) = [a \quad b \quad c \quad d] \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix}$$

Hermite Basis Functions

$$\begin{bmatrix} a & b & c & d \end{bmatrix} \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix} = \begin{bmatrix} h_0 & h_1 & h_2 & h_3 \end{bmatrix} \begin{bmatrix} H_0(t) \\ H_1(t) \\ H_2(t) \\ H_3(t) \end{bmatrix}$$

$$P(t) = \sum_{i=0}^3 h_i H_i(t)$$

Matrix Representation

$$\begin{bmatrix} h_0 \\ h_1 \\ h_2 \\ h_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

Solve for a, b, c, d

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 2 & -2 & 1 & 1 \\ -3 & 3 & -2 & -1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} h_0 \\ h_1 \\ h_2 \\ h_3 \end{bmatrix}$$

Inverse Matrix

Matrix Inverse

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -2 & 1 & 1 \\ -3 & 3 & -2 & -1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

Change Basis

$$\begin{bmatrix} a & b & c & d \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 2 & -3 & 0 & 1 \\ -2 & 3 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix}$$

}

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Change Basis

$$\underbrace{[a \quad b \quad c \quad d] \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix}}_{[h_0 \quad h_1 \quad h_2 \quad h_3]} \begin{bmatrix} 2 & -3 & 0 & 1 \\ -2 & 3 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix}$$

Matrix Transpose

Transpose $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$

$$\left(\begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \right)^T = [a \ b \ c \ d] \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix}$$

Change Basis

$$\underbrace{\begin{bmatrix} a & b & c & d \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix}}_{\begin{bmatrix} h_0 & h_1 & h_2 & h_3 \end{bmatrix}} \underbrace{\begin{bmatrix} 2 & -3 & 0 & 1 \\ -2 & 3 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix}}_{\begin{bmatrix} H_0(t) \\ H_1(t) \\ H_2(t) \\ H_3(t) \end{bmatrix}}$$

Hermite Basis Functions

$$\begin{bmatrix} H_0(t) \\ H_1(t) \\ H_2(t) \\ H_3(t) \end{bmatrix} = \begin{bmatrix} 2 & -3 & 0 & 1 \\ -2 & 3 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} t^3 \\ t^2 \\ t \\ 1 \end{bmatrix}$$


$$H_0(t) = 2t^3 - 3t^2 + 1$$

$$H_1(t) = -2t^3 + 3t^2$$

$$H_2(t) = t^3 - 2t^2 + t$$

$$H_3(t) = t^3 - t^2$$

Hermite Basis Functions



$$H_0(t) = 2t^3 - 3t^2 + 1$$

$$H_1(t) = -2t^3 + 3t^2$$


$$H_2(t) = t^3 - 2t^2 + t$$

$$H_3(t) = t^3 - t^2$$

Ease


A very useful function

In animation, start and stop slowly (zero velocity)




$$H_1(t) = -2t^3 + 3t^2 = t^2(3 - 2t)$$

Catmull-Rom interpolation




**Interpolate points smoothly
Slopes not given though**


Catmull-Rom Interpolation




Catmull-Rom Interpolation



Catmull-Rom Interpolation




Catmull-Rom Interpolation




We can interpolate points as easily as values

Catmull-Rom Interpolation



$$p(t) = (2t^3 - 3t^2 + 1)p_0 + (t^3 - 2t^2 + t)m_0 + (-2t^3 + 3t^2)p_1 + (t^3 - t^2)m_1$$

How to use c-r curve?




N control points
yield
N-1 curve segments

How to choose tangent
condition at two end points?

Video ^_^

- http://v.youku.com/v_show/id_XNTgyNjMwMjM2.html
- 计算机中的数学 (2) — 参变量函数

Bézier curve



Pierre Étienne Bézier
an engineer at Renault



Bézier curve

Bézier curve

$$C(t) = \sum_{i=0}^n P_i B_{i,n}(t), \quad t \in [0,1]$$


where, P_i ($i=0,1,\dots,n$) are control points.

$$B_{i,n}(t) = C_n^i t^i (1-t)^{n-i}, \quad t \in [0,1]$$

Bernstein basis

$$\begin{cases} X(t) = \sum_{i=0}^n x_i B_{i,t}(t) \\ Y(t) = \sum_{i=0}^n y_i B_{i,t}(t) \end{cases}$$

$$C(t) = \begin{pmatrix} X(t) \\ Y(t) \end{pmatrix}, \quad P_i = \begin{pmatrix} x_i \\ y_i \end{pmatrix}$$




Bézier curve


$$\begin{cases} \mathbf{X}(t) = \sum_{i=0}^n x_i B_{i,t}(t) \\ \mathbf{Y}(t) = \sum_{i=0}^n y_i B_{i,t}(t) \end{cases} \quad \begin{cases} \mathbf{X}(t) = \sum_{i=0}^n a_i t^i \\ \mathbf{Y}(t) = \sum_{i=0}^n b_i t^i \end{cases}$$

$$B_{i,n}(t) = C_n^i t^i (1-t)^{n-i}, t \in [0,1]$$


$$C(t) = \begin{pmatrix} \mathbf{X}(t) \\ \mathbf{Y}(t) \end{pmatrix}, \quad P_i = \begin{pmatrix} x_i \\ y_i \end{pmatrix}$$




Bézier curve




(a)




(b)



(c)



(d)



(e)

Bézier curve

Properties of Bernstein basis


$$B_{i,n}(t) = C_n^i t^i (1-t)^{n-i}, t \in [0,1]$$

1. $B_{i,n}(t) \geq 0, i = 0,1,\dots,n, t \in [0,1].$

2. $\sum_{i=0}^n B_{i,n}(t) = 1, t \in [0,1].$

3. $B_{i,n}(t) = B_{n-i,n}(1-t),$
 $i = 0,1,\dots,n, t \in [0,1].$

4. $B_{i,n}(0) = \begin{cases} 1, & i = 0, \\ 0, & \text{else;} \end{cases} B_{i,n}(1) = \begin{cases} 1, & i = n, \\ 0, & \text{else.} \end{cases}$



Bézier curve

Properties of Bernstein basis

5.
$$B_{i,n}(t) = (1-t)B_{i,n-1}(t) + tB_{i-1,n-1}(t), \quad i = 0, 1, \dots, n.$$

6.
$$B'_{i,n}(t) = n[B_{i-1,n-1}(t) - B_{i,n-1}(t)], \quad i = 0, 1, \dots, n.$$

7.
$$(1-t)B_{i,n}(t) = \left(1 - \frac{i}{n+1}\right)B_{i,n+1}(t);$$

$$tB_{i,n}(t) = \frac{i+1}{n+1}B_{i+1,n+1}(t);$$

$$B_{i,n}(t) = \left(1 - \frac{i}{n+1}\right)B_{i,n+1}(t) + \frac{i+1}{n+1}B_{i+1,n+1}(t).$$


Bézier curve

properties of Bézier curves

$$C(t) = \sum_{i=0}^n P_i B_{i,n}(t), \quad t \in [0,1]$$

1. **Endpoint Interpolation:** interpolating two end points

$$C(0) = P_0, \quad C(1) = P_n.$$




2. **tangent direction** of P_0 : P_0P_1 , tangent direction of P_n : $P_{n-1}P_n$.

$$C'(t) = n \sum_{i=0}^{n-1} (P_{i+1} - P_i) B_{i,n-1}(t), \quad t \in [0,1]; \quad C'(0) = n(P_1 - P_0), \quad C'(1) = n(P_n - P_{n-1}).$$

3. **Symmetry:** Let two Bézier curves be generated by ordered Bézier (control) points labelled by $\{p_0, p_1, \dots, p_n\}$ and $\{p_n, p_{n-1}, \dots, p_0\}$ respectively, then the curves corresponding to the two different orderings of control points look the same; they differ only in the direction in which they are traversed.

Bézier curve




3. Symmetry: Let two Bézier curves be generated by ordered Bézier (control) points labelled by $\{p_0, p_1, \dots, p_n\}$ and $\{p_n, p_{n-1}, \dots, p_0\}$ respectively, then the curves corresponding to the two different orderings of control points look the same; they differ only in the direction in which they are traversed.

Bézier curve

properties of Bézier curves

$$C(t) = \sum_{i=0}^n P_i B_{i,n}(t), \quad t \in [0,1]$$



4. Affine Invariance –

the following two procedures yield the same result:


- (1) first, from starting control points $\{p_0, p_1, \dots, p_n\}$ compute the curve and then apply an affine map to it;
- (2) first apply an affine map to the control points $\{p_0, p_1, \dots, p_n\}$ to obtain new control points $\{F(p_0), \dots, F(p_n)\}$ and then find the curve with these new control points.

Bézier curve

properties of Bézier curves

5. **Convex Hull Property** : Bézier curve $C(t)$ lies in the convex hull of the control points P_0, P_1, \dots, P_n ;

6. **variation diminishing property**. Informally this means that the Bezier curve will not "wiggle" any more than the control polygon does..




Bézier curve

Bézier curves

1. linear: $C(t) = (1-t)P_0 + tP_1, t \in [0,1]$,

$$C(t) = [t, 1] \begin{bmatrix} -1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} P_0 \\ P_1 \end{bmatrix}$$



2. quadratic

$$C(t) = (1-t)^2 P_0 + 2t(1-t)P_1 + t^2 P_2$$




$$C(t) = [t^2 \quad t \quad 1] \begin{bmatrix} 1 & -2 & 1 \\ -2 & 2 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} P_0 \\ P_1 \\ P_2 \end{bmatrix}$$

Bézier curve

3. cubic:

$$C(t) = (1-t)^3 P_0 + 3t(1-t)^2 P_1 + 3t^2(1-t)P_2 + t^3 P_3$$

$$C(t) = [t^3 \quad t^2 \quad t \quad 1] \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} P_0 \\ P_1 \\ P_2 \\ P_3 \end{bmatrix}$$



Degree 3


Bézier curve

De Casteljau algorithm

given the control points P_0, P_1, \dots, P_n , and t of Bézier curve, let:

$$P_i^r(t) = (1-t)P_i^{r-1}(t) + tP_{i+1}^{r-1}(t), \quad \forall \left\{ \begin{array}{l} r = 1, \dots, n; \quad i = 0, \dots, n-r \\ P_i^0(u) = P_i \end{array} \right.$$

then $P_0^n(t) = C(t)$.



Bézier curve

Rational Bézier Curve

$$R(t) = \frac{\sum_{i=0}^n B_{i,n}(t)\omega_i P_i}{\sum_{i=0}^n B_{i,n}(t)\omega_i} = \sum_{i=0}^n R_{i,n}(t) P_i$$




Figure 2.19: Circle as Degree 5 Rational Bézier Curve.

where $B_{i,n}(t)$ is Bernstein basis, ω_i is the weight at p_i .

It's a generalization of Bézier curve, which can express more curves, such as circle.

Bézier curve

Properties of rational Bézier curve:

1. endpoints: $R(0) = P_0$; $R(1) = P_n$

2. tangent of endpoints:

$$R'(0) = n \frac{\omega_1}{\omega_0} (P_1 - P_0); \quad R'(1) = n \frac{\omega_{n-1}}{\omega_n} (P_n - P_{n-1})$$

3. Convex Hull Property

.....

5.

6. Influence of the weights




Figure 2.16: Rational Bézier curve.

Bézier surface

Bézier surface

Bézier surface:

$$S(u, v) = \sum_{i=0}^n \sum_{j=0}^m P_{ij} B_{i,n}(u) B_{j,m}(v), \quad 0 \leq u, v \leq 1$$

where $B_{i,n}(u)$ and $B_{j,m}(v)$ Bernstein basis with n degree and m degree, respectively, $(n+1) \times (m+1) P_{i,j} (i=0, 1, \dots, n; j=0, 1, \dots, m)$ construct the control meshes.

